

## Rectal mucosa-associated lymphoid tissue lymphoma treated by endoscopic submucosal resection with band ligation

Katsunori Matsueda, Tatsuya Toyokawa

National Hospital Organization Fukuyama Medical Center, Japan

A 75-year-old woman presented with a positive fecal occult blood test. She had no previous personal or family history of malignancy. Colonoscopy revealed a slightly yellowish 6-mm protrusion in the rectum resembling a submucosal tumor (SMT) (Fig. 1A). Narrow-band imaging showed dilated vessels with branching architectures on the surface and a nonstructural area (Fig. 1B), suggestive of mucosa-associated lymphoid tissue (MALT) lymphoma. However, the color of the lesion was indicative of a carcinoid tumor, which was therefore suspected. Owing to the overlapping features of MALT lymphoma and carcinoid tumor, optical diagnosis of the SMT-like protrusion would be of low confidence. Therefore, it was removed by endoscopic submucosal resection with band ligation (ESMR-L) for diagnosis and treatment (Fig. 1C). Histopathologic examination showed diffuse infiltration of monotonous small-sized lymphoid cells in the mucosa (Fig. 2), positive for CD20 but negative for CD3 and CD10. The lesion was diagnosed as rectal MALT lymphoma. The resected margin was negative. The patient recovered with no complications and remains in complete remission at the five-year follow up.

Whereas ESMR-L is widely used for the treatment of carcinoid tumors [1], our result demonstrates that the technique can also be used for the precise evaluation and safe treatment of a localized, small MALT lymphoma. For an SMT-like protrusion, endoscopic treatment requires special techniques for deeper resection, such as ESMR-L, to achieve clear margins [2]. In the case of a single small SMT-like protrusion in the rectum with low-confidence diagnosis, the lesion should be removed by ESMR-L and sent for precise assessment, difficult with biopsy and endoscopic ultrasound.


Department of Gastroenterology, National Hospital Organization Fukuyama Medical Center, Fukuyama, Japan

Conflict of Interest: None


Correspondence to: Katsunori Matsueda, MD, Department of Gastroenterology, National Hospital Organization Fukuyama Medical Center, 4-14-17 Okinogami-cho, Fukuyama 720-0825, Japan, e-mail: rge43gs@yahoo.co.jp

Received 29 September 2018; accepted 14 October 2018; published online 02 November 2018

DOI: <https://doi.org/10.20524/aog.2018.0324>



**Figure 1** (A) Colonoscopy showed a slightly yellowish, 6-mm submucosal tumor-like protrusion in the rectum. (B) Narrow-band imaging showed dilated vessels with branching architectures on the surface and a nonstructural area. (C) The lesion was removed by endoscopic submucosal resection with band ligation



**Figure 2** Histopathologic examination revealed a mucosa-associated lymphoid tissue lymphoma

### References

1. Bang BW, Park JS, Kim HK, Shin YW, Kwon KS, Kim JM. Endoscopic resection for small rectal neuroendocrine tumors: comparison of endoscopic submucosal resection with band ligation and endoscopic submucosal dissection. *Gastroenterol Res Pract* 2016;**2016**:6198927.
2. Ono A, Fujii T, Saito Y, et al. Endoscopic submucosal resection of rectal carcinoid tumors with a ligation device. *Gastrointest Endosc* 2003;**57**:583-587.