

# Hepatitis C virus genotype 3: clinical features, current and emerging viral inhibitors, future challenges

Vahe Shahnazarian<sup>a</sup>, Daryl Ramai<sup>a,b</sup>, Madhavi Reddy<sup>a</sup>, Smruti Mohanty<sup>c</sup>

The Brooklyn Hospital Center, Academic Affiliate of The Icahn School of Medicine at Mount Sinai, Clinical Affiliate of The Mount Sinai Hospital, Brooklyn, NY; St George's University, Grenada, WI; New York Presbyterian Brooklyn Methodist Hospital, Clinical Affiliate of Weill Cornell Medicine, Brooklyn, NY, USA

## Abstract

Hepatitis C virus (HCV) represents a global burden on healthcare that affects over 150 million people worldwide. In the past, HCV genotype 3 was considered difficult to treat relative to other genotypes. Genotype 3 has been associated with a higher rate of complications, including fatty liver disease, fibrosis, hepatocellular carcinoma and mortality. However, with the advent of first- and second-generation direct-acting antivirals, genotype 3 can be treated effectively. Additionally, these new drugs are well tolerated by patients and have significantly fewer side effects compared to ribavirin and interferon-based regimens. However, while great strides have been made in overcoming biological barriers, our next challenge lies in overcoming economic and financial obstacles if we are to eradicate HCV genotype 3. Herein, we review the clinical features associated with HCV genotype 3, current and emerging treatment regimens, and challenges associated with treatment.

**Keywords** Hepatitis C, genotype 3, sustained viral response, direct acting antivirals, treatment

*Ann Gastroenterol* 2018; 31 (5): 541-551

## Introduction

Viral hepatitis C (HCV) affects approximately 71 million individuals worldwide [1]. A single-stranded RNA virus of the *Flaviviridae* family, HCV is a major cause of chronic liver disease worldwide [2-6]. As a growing public health threat, HCV is one of the leading causes of disability, morbidity and mortality globally [7]. Each year, an estimated 700,000 persons die from HCV-related complications, including

cirrhosis, hepatocellular carcinoma (HCC) and liver failure [8]. Unfortunately, many people with HCV only learn about their infection when they develop symptoms from cirrhosis or liver cancer. It is estimated that 21% of acutely infected patients go undiagnosed, while 75-85% become chronically infected [9]. As a result, there have been renewed international efforts aimed at reducing the global burden of viral hepatitis, including the World Health Organization's (WHO) global health sector strategy on viral hepatitis 2016-2021 [10].

Based on genomic sequence analysis, there are seven major genotypes (HCV 1-7) including 67 confirmed and 20 provisional subtypes [11]. Genotype is not only important in predicting the response to therapy but also predicts the risk of disease progression [12]. Genotype 3 is considered the most poorly understood genotype and has the lowest rate of sustained virologic response (SVR) in the pegylated-interferon era [13]. Regarded as one of the most difficult subtypes to treat, genotype 3 has a higher rate of disease progression and mortality relative to other genotypes [13,14]. According to a meta-analysis, accelerated progression of liver fibrosis associated with genotype 3 had an odds ratio of 1.52 in single biopsy studies compared with other genotypes, suggesting accelerated fibrosis [15]. Moreover, genotype 3 infection has been associated with a higher incidence of HCC [16]. Thus, in the last 20 years, more effective treatments have emerged to combat the virulence of genotype 3.

<sup>a</sup>Division of Gastroenterology, Hepatology, and Advanced Endoscopy, The Brooklyn Hospital Center, Academic Affiliate of The Icahn School of Medicine at Mount Sinai, Clinical Affiliate of The Mount Sinai Hospital, Brooklyn, NY (Vahe Shahnazarian, Daryl Ramai, Madhavi Reddy); <sup>b</sup>School of Medicine, St George's University, True Blue, Grenada, WI (Daryl Ramai); <sup>c</sup>Division of Gastroenterology and Hepatology, New York Presbyterian Brooklyn Methodist Hospital, Clinical Affiliate of Weill Cornell Medicine, Brooklyn, NY (Smruti Mohanty), USA

Conflict of Interest: None

Correspondence to: Vahe Shahnazarian, MD, MPH, Gastroenterology and Hepatology Fellow, Division of Gastroenterology, Hepatology, and Advanced Endoscopy, The Brooklyn Hospital Center, Academic Affiliate of The Icahn School of Medicine at Mount Sinai, Clinical Affiliate of The Mount Sinai Hospital, Brooklyn, NY 11201, USA, e-mail: vaheshahnazarian@gmail.com

Received 12 February 2018; accepted 18 April 2018; published online 4 June 2018

DOI: <https://doi.org/10.20524/aog.2018.0281>

## Methodology

Studies published between January 2000 and January 2018 were searched in electronic databases according to article titles, abstract contents, and relevance in the field of viral hepatitis C genotype 3. The databases used in this review included PubMed, MEDLINE, Embase, Google Scholar, Scopus, ClinicalTrials.gov and Cochrane Databases. The main search terms used were hepatitis C genotype 3, epidemiology, infection rates, steatosis, fibrosis, HIV coinfection, HCC, insulin resistance (IR), lipid metabolism, treatment, and clinical trials. We also manually searched references to identify additional relevant studies. Articles published in languages other than English were excluded.

## Clinical features

### Epidemiology

Genotype 3 represents 20% of all HCV infections worldwide [17]. It is the second most common genotype in the world, affecting approximately 54.3 million individuals, with approximately 75% of cases occurring in East Asia [18]. Genotype 3 is responsible for 54-80% of all HCV infections in India, and 79% of all HCV infections in Pakistan, with higher rates reported among drug users [17,19]. It is also prevalent in Europe where it accounts for 30% of all HCV infections in Greece, Poland, Denmark, Finland and the UK [19]. In the US, genotype 3 is the second most common genotype and accounts for 8-13% of all HCV infections [20]. In Africa, genotype 3 is far less prevalent and makes up approximately 9% of all HCV infections [21].

### Steatosis

Steatosis occurs as a result of the excessive accumulation of triglycerides within hepatocytes. A study reviewing liver histopathology found that genotype 3 had the highest prevalence of steatosis when compared to other subtypes [22]. This is likely due to the cytopathic effect of the genotype, which selectively interferes with the late cholesterol pathway and upregulates the expression of fatty acid synthase; however, steatosis tends to improve with SVR [23,24]. Steatosis has also been noted to progress more rapidly into liver fibrosis in patients infected with genotype 3 rather than other genotypes [25]. However, Chan *et al* reported that while genotype 3 is associated with higher rates of steatosis, current data do not show any enhanced direct viral-mediated hepatic fibrogenesis [21]. Most studies that reported a potential association with genotype 3 and severity of hepatic fibrosis were limited by small cohorts, variability in patient characteristics, such as insulin resistance (IR), body mass index or genotype distribution, and discrepancies in methodology, particularly with respect to the grading of steatosis [26,27].

A large meta-analysis of 3068 chronic hepatitis C patients from North America, Europe and Australia showed that genotype 3 was associated with steatosis, not fibrosis; multivariate analysis identified steatosis and the level of inflammatory activity on histopathology as independent predictors of disease [28]. While a direct viral association seems lacking or unclear, some studies have identified a link between higher grades of steatosis and higher rates of fibrosis [27].

While the precise pathogenic mechanisms of genotype 3-mediated steatosis are still unclear, genotype 3 appears to modulate host lipid metabolism and transportation within hepatocytes via three mechanisms: microsomal triglyceride transfer protein; sterol regulatory element-binding protein 1c; and peroxisome proliferator-associated receptor- $\alpha$  [29]. Through these pathways, it is thought that intrahepatic fat sequestration or steatosis impairs the response to treatment in patients with fatty liver [25]. In a large national study of 7248 eligible subjects, the use of statins (n=3334) was significantly associated with better SVR rates after antiviral therapy and lower progression of liver fibrosis and incidence of HCC [30].

### HCV/human immunodeficiency virus (HIV) coinfection

Coinfection with HIV and HCV-3 is relatively common. According to a meta-analysis of over 780 studies, the global prevalence of HCV/HIV coinfection was estimated to be about 6% [31]. In the US and Western Europe, the prevalence of coinfection is approximately 10-30% [32,33]. Coinfection with HCV-3 increases the risk of chronicity while producing a higher viral load and an inability to mount a CD4/CD8 mediated T-cell immune response [34]. Coinfection has also been associated with an accelerated progression of liver fibrosis and a reduced SVR to interferon-based regimens [35-37]. However, antiretroviral therapy has been shown to ease the progression of HCV-associated liver injury and fibrosis by reducing HIV-related inflammation and immune dysfunction, and limiting infectivity [38-41]. Additionally, HCV treatment in coinfecting patients has been associated with a reduction in the frequency of death, HIV progression, liver-related events, and reduced hazards of diabetes mellitus and possibly chronic renal failure [42,43]. Since HIV accelerates the natural history of HCV and liver-related complications, it is recommended that all coinfecting patients be treated for chronic HCV, except those whose life expectancy is less than a few years and will not be remediated by treatment [44].

### Hyperlipidemia, diabetes mellitus or IR

Lipid metabolism is intimately involved in the molecular mechanisms of the HCV infectious cycle. HCV replication influences and depends upon cholesterol uptake and efflux through different lipoprotein receptors during its entry into the host's cells [45]. Very low-density lipoprotein-associated proteins, including apolipoprotein (apo)B, apoE and

microsomal triglyceride transfer proteins, have been shown to play a crucial role in the formation of infectious HCV particles, especially pertaining to genotype 3 [46]. HCV can bind low-density lipoprotein (LDL) receptors and lead to intracellular lipid deposition [47]. Patients infected with genotype 3 tend to have hypocholesterolemia and hypobetalipoproteinemia, which may account for the direct effect of the virus on lipid metabolism [48]. It appears that HCV-3 also selectively interferes with the late cholesterol pathway, a phenomenon that appears to disappear with SVR [23]. Interestingly, high LDL levels tends to predict SVR in patients treated with interferon and ribavirin (RBV) [49]. Since LDL receptors are involved in HCV entry into hepatocytes, higher LDL levels may decrease the number of LDL receptors on the cell membrane, thus decreasing cellular infectivity [49].

HCV core protein can induce IR by disturbing the intracellular signaling pathway of insulin [50]. In genotype 3, the presence of IR may reduce the probability of achieving SVR using interferon and RBV regimens, especially in the setting of steatosis involving more than 33% of the liver [51-54]. However, eradication of HCV appears to improve insulin sensitivity without altering fat deposits, adipokine or glucagon levels, suggesting a direct link between viral activity and IR [55]. Subjects with normal insulin sensitivity were almost three times more likely to attain SVR than those with IR [56]. Unfortunately, a prospective cohort study was unable to find an association between IR and SVR rates in genotype 1, 2 or 3 patients [57]. Additionally, Patel *et al* looked at genotype-specific associations with IR and liver fibrosis. While the study did report an association in genotype 1 patients, no significant association was found in the genotype 3 cohort [58]. The mechanisms of HCV-mediated IR, SVR, and the genotype-specific associations remain unclear. Fortunately, new direct-acting antiviral (DAA) regimens appear unaffected by IR [54].

## HCC

The incidence of HCC has been on the rise in the United States and around the world [59,60]. Genotype 3 has been associated with significantly higher rates of HCC relative to other genotypes [61]. McMahon *et al* studied 1080 participants for 11,171 person-years and reported that individuals infected with genotype 3 were at increased risk for HCC and liver-related death [62]. Another study showed that the risk of HCC, after adjusting for demographic and clinical characteristics, was 80% higher in patients with genotype 3 when compared to genotype 1 [63]. Likewise, McCombs *et al* showed in a Veteran Affairs (VA) study that the risk for developing HCC was 63% higher with genotype 3 (HR 1.63, 95%CI 1.47-1.79;  $P < 0.001$ ) compared with genotype 1 [14]. In addition, Ioannou *et al* (2018) reported that in 62,354 VA patients the treatment of HCV with DAAs reduced the risk of liver cancer by 71% [64]. The finding emphasizes the need to be vigilant in the diagnosis and early treatment of HCV even with cirrhosis of the liver.


## Current DAAs

The genetic landscape exhibited by HCV has created a major challenge for developing vaccines and effective pan-genotypic therapies [65]. Prior to DAAs, pegylated interferon and RBV regimens were the standard of care. However, the main limitation of those regimens was their side effects, dominated by fatigue, influenza-like symptoms, hematologic abnormalities, and neuropsychiatric symptoms [66]. These side effects have been subverted with DAAs, which treat patients facing major challenges including HCV genetic diversity, viral resistance, the influence of host genetics, advanced liver disease and other comorbidities [67-69].

Major advances in elucidating and characterizing the life cycle of HCV have led to the targeting of key stages of viral replication [70,71] (Fig. 1). DAAs target non-structural (NS) viral proteins crucial to viral RNA replication, polyprotein processing, and the assembly of infectious virions [72]. Currently, DAAs approved by the FDA for the treatment of genotype 3 are (in combination): sofosbuvir, a pangenotypic nucleotide analog inhibitor of HCV NS5B polymerase; daclatasvir, an inhibitor of HCV NS5A metalloprotein; velpatasvir, a next-generation pangenotypic NS5A inhibitor; and glecaprevir and pibrentasvir, a NS3/4A protease inhibitor and NS5A inhibitor, respectively [73,74].

The ASTRAL-3 trials aimed to evaluate the safety and efficacy of sofosbuvir and velpatasvir in the treatment of patients infected with genotype 3, including treatment-naïve, treatment-experienced patients, and patients with compensated liver cirrhosis [75]. After 12-week treatment, the sofosbuvir-velpatesvir group achieved an SVR of 95% (95%CI 92-98%), superior to the rate of 80% (95%CI 75-85%) in the sofosbuvir-RBV group ( $P < 0.001$ ). The most frequently reported side effects were fatigue, headache, nausea, and insomnia.

The combination of glecaprevir and pibrentasvir was tested in the SURVEYOR-2 and ENDURANCE-3 trials [76]. In SURVEYOR-2 phase II, the combination was tested with or without RBV. The study reported an SVR of 97% for treatment-naïve genotype 3 patients without cirrhosis after 8 weeks of treatment (without RBV). Following 12 weeks of therapy (with or without RBV), treatment-naïve genotype 3 patients showed an SVR of 100%. In SURVEYOR-2 phase III, treatment-experienced patients had either 12 or 16 weeks of glecaprevir/pibrentasvir; in those with cirrhosis treatment-naïve patients had 12 weeks and treatment-experienced patients 16 weeks of therapy [77]. In treatment-experienced, non-cirrhotic patients, cure rates of 91% and 96% were reported in the 12- and 16-week cohorts, respectively. In the cirrhotic arms, treatment-naïve patients had an SVR of 98% after the 12-week regimen while the treatment-experienced patients had an SVR of 96% after 16 weeks. Studies comparing the efficacy of glecaprevir/pibrentasvir (for 8 weeks) and sofosbuvir/daclatasvir in treatment-naïve non-cirrhotics reported similar safety and SVR profiles [78]. The study suggested that an 8-week treatment of glecaprevir/pibrentasvir could prove highly efficacious. Adding to the guidelines issued by the European Association for the Study of the Liver (EASL), the American Association for the



**Figure 1** Illustration of stages at which NS5A and NS5B inhibitors take effect

Study of Liver Diseases and the Infectious Disease Society of America recommend the use of glecaprevir/pibrentasvir for 8 weeks in individuals without cirrhosis (Table 1) [79].

In the PROTON trial, a combination of sofosbuvir, interferon and RBV was administered for 12 weeks. It showed an SVR of 96% in treatment-naïve genotype 3 patients, some of whom had cirrhosis [80]. The BOSON trial, a phase 3 trial, evaluated the same combination, with an SVR of 93% in their treatment-naïve patients (96% in non-cirrhotics and 91% in cirrhotics) [78]. Interferon-free cohorts receiving sofosbuvir and RBV experienced lower SVRs of 71% and 84% after being treated for 16 and 24 weeks, respectively.

In the FISSION trial, sofosbuvir and RBV for 12 weeks was compared to interferon and RBV for 24 weeks in treatment-naïve patients [81]. However, the SVRs were only 56% and 63%, respectively. In both groups, cirrhotics had a lower SVR than those without cirrhosis (34% vs. 61% and 30% vs. 71%, respectively). The POSITRON trial was for patients in whom interferon was not an option [82]. Subjects received sofosbuvir and RBV or placebo for 12 weeks and reported SVRs of 61% and 0% respectively. Cirrhotics had an SVR of 21% which non-cirrhotics achieved an SVR of 68%. Likewise, the VALENCE trial extended the same drug combination of sofosbuvir and RBV to 24 weeks in treatment-naïve patients. However, a much higher SVR was seen in treated patients without cirrhosis (94%) and in cirrhotics (92%) [83].

The ALLY-3 study included treatment-naïve patients who received a combination of sofosbuvir and daclatasvir over 12 weeks. The combination achieved an SVR of 90% and was well tolerated; the SVR was 58% in cirrhotics and 97% in non-cirrhotics [84,85]. ALLY-3 also tested treatment-experienced

patients with the same regimen and reported an SVR of 86% overall (69% in cirrhotics and 94% in non-cirrhotics) (Table 2).

Velpatasvir (GS-5816) is a potent NS5A inhibitor with activity against genotypes 1-6 [86]. In the ELECTRON-2 trial, it was evaluated at different doses in combination with different drugs: sofosbuvir with 25 mg of velpatasvir, sofosbuvir with 25 mg of velpatasvir and RBV, sofosbuvir with 100 mg of velpatasvir, and sofosbuvir with 10 mg of velpatasvir and RBV, all in treatment-naïve patients [87]. The study reported SVRs of 100%, 88%, 96% and 100%, respectively. It was also tested in treatment-experienced patients in the same four arms, where SVRs were 85%, 95%, 100% and 100% in non-cirrhotic patients, and 58%, 85%, 88% and 96% in cirrhotic patients, respectively. The most common adverse effects were fatigue, headache and nausea, all related to RBV.

A recent meta-analysis by Berden *et al* showed that regimens containing sofosbuvir and velpatasvir achieved the highest treatment efficacy with RBV (SVR 99%; 95%CI 98-100%) and without RBV (SVR 97%; 95%CI 95-99%) [88]. Analysis showed that RBV significantly increased SVR rates in patients with or without liver cirrhosis. The same regimen has also demonstrated high treatment efficacy (SVR 94%) in patients coinfecting with HIV-1 and HCV [89]. As a result, according to the EASL recommendations for the treatment of HCV, genotype 3 mono-infected or HIV-coinfecting patients should be treated with sofosbuvir and daclatasvir or with sofosbuvir and velpatasvir (for 12 weeks in treatment-naïve individuals) [90]. Treatment-experienced individuals are recommended either combinations for an extended duration of 24 weeks or the addition of RBV for 12 weeks.

**Table 1** American Association for the Study of Liver Diseases: genotype 3 treatment recommendations

Population	Cirrhosis status	Recommendation	Combination	Duration	SVR
Treatment-naïve	Without cirrhosis	1 <sup>st</sup> line	G/P SOF/VEL	12 wk 8 wk	95% 98%
		2 <sup>nd</sup> line	DCV PO daily + SOF w/ or w/o RBV	12 wk	90%
	With compensated cirrhosis	1 <sup>st</sup> line	SOF/VEL G/P	12 wk 12 wk	93% 100%
		2 <sup>nd</sup> line	DCV + SOF w/ or w/o weight based RBV	24 wk	86%
			SOF/VEL/voxilaprevir*	12 wk	96%
		Treatment-experienced	Without cirrhosis	1 <sup>st</sup> line	SOF/VEL
2 <sup>nd</sup> line	DCV + SOF			12 wk	94%
	G/P			16 wk	96%
	SOF/VEL/voxilaprevir			12wk	84%
With Compensated Cirrhosis	1 <sup>st</sup> line		SOF/VEL/voxilaprevir Elbasvir/grazoprevir + SOF	12 wk 12 wk	96% 100%
	2 <sup>nd</sup> line		SOF/VEL w/ weight based RBV G/P	12 wk 16 wk	89% 96%
Treatment-naïve	With decompensated cirrhosis and RBV eligible	1 <sup>st</sup> line	DCV + SOF w/ low initial dose of RBV (increase as tolerated)	12 wk	83-94%
			SOF/VEL w/ weight based RBV	12 wk	85%
Treatment-naïve	With decompensated cirrhosis and RBV ineligible	1 <sup>st</sup> line	DCV + SOF	24 wk	60%
			SOF/VEL	24 wk	50%
Treatment-experienced	With decompensated cirrhosis and Sofosbuvir- or NS5A-based treatment failure	1 <sup>st</sup> line	SOF/ VEL	24 wk	78%

\*When RAS Y93H is present

SVR, sustained virologic response; SOF, sofosbuvir; IFN, interferon; RBV, ribavirin; DCV, daclatasvir; G, glecaprevir; P, pibrentasvir; VEL, velpatasvir

Today, many of these treatments are very expensive compared to prior regimens of interferon and RBV. The sofosbuvir, ledipasvir and RBV regimen costs approximately \$73,000 per quality-adjusted life year (QALY), whereas the sofosbuvir and daclatasvir combination costs more than \$396,000 per QALY [19]. However, from a societal perspective, treatment with novel HCV therapies appears to be cost-effective compared to RBV and pegylated interferon-based treatments [91,92]. The use of DAAs has led to dramatically higher SVR rates along with fewer side effects. The current recommendations for the treatment of chronic HCV-genotype 3 are summarized in Table 1.

### Emerging next-generation antivirals

Future developments consist of next-generation DAAs capable of overcoming resistance barriers and offer a broader spectrum of activity against all genotypes, including glecaprevir, pibrentasvir, voxilaprevir, uprifosbuvir, grazoprevir, ruzasvir, AL-335, simeprevir, and odalasvir [93]. The C-SWIFT trial combined grazoprevir, elbasvir and sofosbuvir in the

treatment of naïve patients with and without cirrhosis [94]. Non-cirrhotic patients received the therapy for 8 or 12 weeks and achieved SVRs of 93% and 100%, respectively. Cirrhotic patients achieved an SVR of 91% after a 12-week course. Similarly, the C-ISLE study assessed the efficacy and safety of elbasvir and grazoprevir plus sofosbuvir with and without RBV, though in patients with compensated cirrhosis [95]. Among treatment-naïve participants, SVR at 8-weeks was 91% (21/23) in those treated with RBV. Following treatment for 12-weeks without RBV, SVR was 96% (23/24). Among treatment-experienced participants, with and without RBV, SVR was 94% (17/18) and 100% (17/17) in the 12-week arm, respectively, and 94% (17/18) in the 16-week arm. Five individuals reported significant adverse effects, including pneumonia, chest pain, opiate overdose, cellulitis, and decreased creatinine. Paritaprevir (ABT-450), an NS3/4 inhibitor, is administered with ritonavir (ABT-450/r) [96]. As a pharmaco-enhancer, ritonavir has no activity against HCV; instead, it inhibits the metabolism of paritaprevir, thereby increasing peak and trough drug exposures and allowing for a once-daily dose of ABT-450 [97]. The NAVIGATOR trial combined ABT-450/r with ombitasvir (ABT-267), an NS5A

inhibitor, for 12 weeks in genotype 3 treatment-naïve non-cirrhotic patients, achieving SVR rates of 50% without RBV and just 9% with RBV [98]. Though the medication was well tolerated, the low SVRs prompted discontinuation of the trial. Similarly, in patients with or without cirrhosis, the QUARTZ II-III trial combinations of ombitasvir/paritaprevir/ritonavir plus sofosbuvir with or without RBV showed an overall SVR of 98% after 12 weeks [98]. The study noted that combining DAAs with complementary mechanisms of action and virological targets may be an effective therapeutic strategy.

Other novel inhibitors are being studied, including uprifosbuvir (MK-3682; NS5B inhibitor), ruzasvir (MK8408; NS5A inhibitor), grazoprevir (NS3 inhibitor), and voxilaprevir (NS3/4A inhibitor) [99]. The C-BREEZE-2 trial looked at the efficacy of ruzasvir and uprifosbuvir in treating genotypes 1 to 6 for 12 weeks [100]. However, because of the suboptimal genotype 3 SVRs (80% in patients without cirrhosis vs. 68% in patients with cirrhosis), as well as virological relapse, the combination was deemed therapeutically ineffective (Table 3).

In phase II of C-CREST-2, a triple combination of uprifosbuvir, ruzasvir and grazoprevir was tested [101]. The overall SVR in genotype 3, treatment-naïve, non-cirrhotic patients ranged from 86-95%, suggesting that the 8-week treatment duration was suboptimal [102]. In phase III of the C-CREST 2 trial, the same triple combination, with and without RBV, was investigated at 8, 12, and 16 weeks [102]. The overall SVR for genotype 3 patients was 96%. In the 8-week arm, SVR was 94% and 98%, without and with RBV, respectively. In the cirrhotic subgroup, only 1 patient relapsed out of 74, and this patient was in the 16-week treatment (with RBV) group.

Phase II of the LEPTON trial evaluated a triple combination of sofosbuvir, velpatasvir and GS-9857, a next-generation NS3/4A protease inhibitor [100]. The treatment was tested in treatment-naïve patients with compensated cirrhosis, interferon/RBV failures with cirrhosis, and DAA failures with and without cirrhosis. The naïve group had a duration of 6 weeks while the experienced group had a duration of 8 weeks. The naïve group had an SVR of 83% while the experienced group had an SVR of 100%, which included four prior DAA non-responders. However, the study was inadequately powered and these results should therefore be interpreted cautiously.

In the POLARIS-3 trial, treatment-naïve and experienced patients with cirrhosis were treated with sofosbuvir and velpatasvir for 12 weeks, or sofosbuvir, velpatasvir, and voxilaprevir for 8 weeks [103]. The trial enrolled patients who had not previously received treatment with DAAs. Overall, both groups achieved an SVR of 96% after 12-weeks, which included one withdrawn consent and an unrelated death. Treatment-naïve patients with cirrhosis had an SVR of 96% on the triple therapy, while treatment-experienced patients with cirrhosis had an SVR of 97%. About 2% of study participants experienced adverse events.

Finally, MIV-802, an NS5B inhibitor, was tested to assess its antiviral activity against genotype 3 *in vitro* [104]. Preclinical studies found that MIV-802 had greater antiviral activity against genotype 3 than did sofosbuvir,

which supported its consideration for further development [105]. Overall, next-generation HCV DAAs appear to be clinically effective in treating a wide patient population. SVR rates for next-generation DAAs are either equal to or higher than those for first-generation DAAs. This new class of therapeutics promise to be easy to use, highly efficient, pan-genotypic, once-daily, all-oral, and interferon- and RBV-free [93].

### Future challenges

HCV drug development in the last decade has overcome many challenges and has addressed gaps in care, such as pan-genotype coverage, adverse side effects, and coverage for HIV coinfecting individuals and cirrhotics. Achieving clinical SVR has become less of an issue in genotype 3 infected patients, once considered difficult to treat [106]. Studies in certain patient populations, such as children, pregnant women and nursing mothers, are lacking and represent an unmet need [107]. Additionally, while most HCV medications are metabolized by the liver, DAA options for persons with severe chronic kidney disease remain limited or contraindicated [108]. Furthermore, the sample size from subgroup analysis emerging from the clinical trials of next-generation DAAs must be interpreted cautiously. As a result, larger sample sizes are needed to evaluate the clinical efficacy within special cohorts.

Moreover, given that cost plays a role in the distribution and uptake of treatments, it remains to be seen how cost will affect public health. Our next challenge lies in overcoming cost barriers if we are to truly eradicate HCV genotype 3 [109]. We must address drug prices to ensure universal access. We must also combine these efforts with harm reduction aimed at reintroducing new and resistant strains of HCV. More importantly, in order to achieve the strategic goal set by the WHO, which aims to eliminate HCV, researchers, healthcare providers, policy makers, affected communities, advocates, pharmaceutical and diagnostics industries, and governments around the world will have to work together [110]. Falade-Nwulia *et al* noted that reducing reinfection and achieving HCV elimination goals involve coupling HCV treatment with harm reduction and targeted behavioral interventions; reinfection rates should be viewed as an indicator of the uptake of HCV treatment [111].

Furthermore, although DAAs have been successful in treating HCV, their financial burden does not exclude the possibility of the use and development of a vaccine that may serve a crucial preventative role [112]. While vaccines are available for preventing hepatitis A and B, the goal of HCV vaccine development is protection from HCV persistence. Several candidate vaccines have progressed to clinical testing after promising results in chimpanzees [113]. Two vaccination mechanisms are currently being investigated using T-cell mediated immunity and eliciting antibodies [114]. However, the complex transcriptional profile of hepatitis C continues to be challenge vaccine development.

**Table 2** Current treatment options with direct acting antivirals

Population	Study	Combination	SVR (%)
Treatment-naïve	PROTON	SOF + IFN + RBV 12 wk	No cirrhosis: 96
	FISSION	SOF + RBV 12 wk	No cirrhosis: 61
			Cirrhosis: 34
	POSITRON	SOF + RBV 12 wk	No cirrhosis: 68
	VALENCE		Cirrhosis: 21
		SOF + RBV 24 wk	No Cirrhosis: 94
	BOSON		Cirrhosis: 92
		SOF + IFN + RBV 12 wk	No cirrhosis: 96
			Cirrhosis: 91
		SOF + RBV 16 wk	No cirrhosis: 83
			Cirrhosis: 57
		SOF + RBV 24 wk	No cirrhosis: 90
		Cirrhosis: 82	
	ALLY-3	SOF + DCV 12 wk	No cirrhosis: 97
	Cirrhosis: 58		
ELECTRON-2	SOF + LDV 12 wk	No cirrhosis: 64	
	SOF + LDV + RBV 12 wk	No cirrhosis: 100	
SURVEYOR-2	G/P + RBV 8 wk	No cirrhosis: 97	
	G/P + RBV 12 wk	Cirrhosis: 100	
ASTRAL-3	SOF + VEL 8 wk	No cirrhosis: 100	
	SOF + VEL + RBV 8 wk	No cirrhosis: 88	
	SOF + VEL 8 wk	No cirrhosis: 96	
	SOF + VEL + RBV 8 wk	No cirrhosis: 100	
	SOF + VEL 12 wk	Cirrhosis: 91	
Treatment-experienced	LONESTAR-2	SOF + IFN + RBV 12 wk	No cirrhosis: 83
		Cirrhosis: 83	
	FUSION	SOF + RBV 12 wk	No cirrhosis: 37
			Cirrhosis: 19
		SOF + RBV 16 wk	No cirrhosis: 63
			Cirrhosis: 61
	VALENCE	SOF + RBV 24 wk	Overall: 70
			No cirrhosis: 85
			Cirrhosis: 60
	BOSON	SOF + IFN + RBV 12 wk	No cirrhosis: 94
			Cirrhosis: 86
		SOF + RBV 16 wk	No cirrhosis: 76
		Cirrhosis: 47	
		SOF + RBV 24 wk	No cirrhosis: 82
		Cirrhosis: 77	
ELECTRON-2		SOF + LDV + RBV 12 wk	No cirrhosis: 89
		Cirrhosis: 73	
SURVEYOR-2	G/P + RBV 12 wk	No cirrhosis: 91	
	G/P + RBV 16 wk	No cirrhosis: 96	
	G/P + RBV 12 wk	Cirrhosis: 96	

SVR, sustained virologic response; SOF, sofosbuvir; IFN, interferon; RBV, ribavirin; DCV, daclatasvir; LDV, ledipasvir; VEL, velpatasvir; G/P, glecaprevir / pibrentasvir

HCV reinfection should be minimized but not feared as we work towards HCV elimination. Rather, reinfection should be viewed as an indication of the uptake of hepatitis C treatment in populations most likely to sustain ongoing HCV epidemics. HCV reinfections should be minimized by coupling HCV treatment with harm reduction and well-designed targeted behavioral interventions.

### Concluding remarks

HCV represents a global health threat. Specifically, genotype 3 is known to have a rapid disease progression relative to other genotypes, leading to advanced stages of liver disease far earlier. Through the manipulation of lipid

**Table 3** Next-generation treatments

Population	Study	Combination	SVR (%)
Treatment-naïve	C-SWIFT	Elbasvir + Grazoprevir + SOF	No cirrhosis, 8 wk: 93 No cirrhosis, 12 wk: 100 Cirrhosis, 12 wk: 91
	C-ISLE	Elbasvir + Grazoprevir + SOF	Cirrhosis, 8 wk: 91 Cirrhosis, 12 wk: 96
	NAVIGATOR	ABT-450/r + Ombitasvir ABT-450/r + Ombitasvir + RBV	No cirrhosis, 12 wk: 40 No cirrhosis, 12 wk: 9
	QUARTZ II-III	Ombitasvir + Paritaprevir + Ritonavir + SOF Ombitasvir + Paritaprevir + Ritonavir + SOF + RBV Ombitasvir + Paritaprevir + Ritonavir + SOF + RBV	No cirrhosis, 12 wk: 98 No cirrhosis, 12 wk: 91 Cirrhosis, 12 wk: 100
	C-CREST-2	Ruzasvir + Uprifosbuvir + Grazoprevir Ruzasvir + Uprifosbuvir + Grazoprevir + RBV Ruzasvir + Uprifosbuvir + Grazoprevir Ruzasvir + Uprifosbuvir + Grazoprevir + RBV	No cirrhosis, 8/12 wk: 95 No cirrhosis, 12 wk: 100 Cirrhosis, 12 wk: 97 Cirrhosis, 12 wk: 100
	LEPTON	SOF + VEL + GS-9857	Cirrhosis, 6 wk: 83 Cirrhosis, 8 wk: 100
	POLARIS-3	SOF + VEL + Voxilaprevir SOF + VEL	Cirrhosis, 8 wk: 96 Cirrhosis, 12 wk: 96
Treatment-experienced	C-ISLE	Elbasvir + Grazoprevir + SOF	Cirrhosis, 12 wk: 100 Cirrhosis, 16 wk: 94
	LEPTON	SOF + VEL + GS-9857	No cirrhosis, 8 wk: 100 Cirrhosis, 8 wk: 100
	C-CREST-2	Ruzasvir + Uprifosbuvir + Grazoprevir Ruzasvir + Uprifosbuvir + Grazoprevir + RBV	No cirrhosis, 12 wk: 97 Cirrhosis, 12 wk: 100 No cirrhosis, 12 wk: 93 Cirrhosis, 12 wk: 100

SVR, sustained virologic response; SOF, sofosbuvir; IFN, interferon; RBV, ribavirin; VEL, velpatasvir; ABT-450/r, Paritaprevir/ritonavir

metabolic pathways, steatosis impairs the treatment response. However, even in patients with advanced liver disease, new DAA regimens are slowly turning the tide against genotype 3. A high SVR can be achieved in patients with a spectrum of liver disease, as well as those coinfecting with HIV. New DAA regimens are also better tolerated by patients, since they have fewer adverse side-effects. We believe that with this expanded armamentarium, genotype 3 should no longer be considered difficult to treat.

## References

1. Polaris Observatory HCV Collaborators. Global prevalence and genotype distribution of hepatitis C virus infection in 2015: a modelling study. *Lancet Gastroenterol Hepatol* 2017;**2**:161-176.
2. Lavanchy D. The global burden of hepatitis C. *Liver Int* 2009;**29**(Suppl 1):74-81.
3. Chen SL, Morgan TR. The natural history of hepatitis C virus (HCV) infection. *Int J Med Sci* 2006;**3**:47-52.
4. Seeff LB. The history of the “natural history” of hepatitis C (1968-2009). *Liver Int* 2009;**29**(Suppl 1):89-99.
5. Kleven RM, Hu DJ, Jiles R, Holmberg SD. Evolving epidemiology of hepatitis C virus in the United States. *Clin Infect Dis* 2012;**55**(Suppl 1):S3-S9.
6. Smith BD, Morgan RL, Beckett GA, Centers for Disease Control and Prevention. Recommendations for the identification of chronic hepatitis C virus infection among persons born during 1945-1965. *MMWR Recomm Rep* 2012;**61**:1-32.
7. Stanaway JD, Flaxman AD, Naghavi M, et al. The global burden of viral hepatitis from 1990 to 2013: findings from the Global Burden of Disease Study 2013. *Lancet* 2016;**388**:1081-1088.
8. World Health Organization. WHO issuing updated guidelines for treatment of hepatitis C infection. <http://www.who.int/hepatitis/news-events/hepatitis-c-guidelines-2016-story/en/> [Accessed May 1, 2018]
9. Centers for Disease Control and Prevention. The ABCs of hepatitis. <http://www.cdc.gov/hepatitis/Resources/Professionals/PDFs/ABCTable.pdf> [Accessed Jul 28, 2018]
10. WHO. Global health sector strategy on viral hepatitis 2016-2021. Towards ending viral hepatitis. <http://apps.who.int/iris/bitstream/10665/246177/1/WHO-HIV-2016.06-eng.pdf> [Accessed Jul 28, 2018]
11. Smith DB, Bukh J, Kuiken C, et al. Expanded classification of hepatitis C virus into 7 genotypes and 67 subtypes: updated criteria and genotype assignment web resource. *Hepatology* 2014;**59**:318-327.
12. Ge D, Fellay J, Thompson AJ, et al. Genetic variation in IL28B predicts hepatitis C treatment-induced viral clearance. *Nature* 2009;**461**:399-401.
13. Sarin SK, Kumar CK. Treatment of patients with genotype 3 chronic hepatitis C—current and future therapies. *Liver Int* 2012;**32**(Suppl 1):141-145.



14. McCombs J, Matsuda T, Tonnu-Mihara I, et al. The risk of long-term morbidity and mortality in patients with chronic hepatitis C: results from an analysis of data from a Department of Veterans Affairs Clinical Registry. *JAMA Intern Med* 2014;**174**:204-212.
15. Probst A, Dang T, Bochud M, Egger M, Negro F, Bochud PY. Role of hepatitis C virus genotype 3 in liver fibrosis progression—a systematic review and meta-analysis. *J Viral Hepat* 2011;**18**:745-759.
16. Nkontchou G, Zioli M, Aout M, et al. HCV genotype 3 is associated with a higher hepatocellular carcinoma incidence in patients with ongoing viral C cirrhosis. *J Viral Hepat* 2011;**18**:e516-e522.
17. Gower E, Estes C, Blach S, Razavi-Shearer K, Razavi H. Global epidemiology and genotype distribution of the hepatitis C virus infection. *J Hepatol* 2014;**61**:S45-S57.
18. Messina JP, Humphreys I, Flaxman A, et al. Global distribution and prevalence of hepatitis C virus genotypes. *Hepatology* 2015;**61**:77-87.
19. Ampuero J, Romero-Gómez M, Reddy KR. Review article: HCV genotype 3 – the new treatment challenge. *Aliment Pharmacol Ther* 2014;**39**:686-698.
20. Kohli A, Shaffer A, Sherman A, Kotttilil S. Treatment of hepatitis C: a systematic review. *JAMA* 2014;**312**:631-640.
21. Chan A, Patel K, Naggie S. Genotype 3 infection: the last stand of hepatitis C virus. *Drugs* 2017;**77**:131-144.
22. Gondeau C, Pageaux GP, Larrey D. Hepatitis C virus infection: Are there still specific problems with genotype 3? *World J Gastroenterol* 2015;**21**:12101-12113.
23. Clark PJ, Thompson AJ, Vock DM, et al. Hepatitis C virus selectively perturbs the distal cholesterol synthesis pathway in a genotype-specific manner. *Hepatology* 2012;**56**:49-56.
24. Lee YA, Friedman SL. Reversal, maintenance or progression: what happens to the liver after a virologic cure of hepatitis C? *Antiviral Res* 2014;**107**:23-30.
25. Buti M, Esteban R. Hepatitis C virus genotype 3: a genotype that is not 'easy to treat'. *Expert Rev Gastroenterol Hepatol* 2014;**9**:375-385.
26. Westin J, Nordlinder H, Lagging M, Norkrans G, Wejstål R. Steatosis accelerates fibrosis development over time in hepatitis C virus genotype 3 infected patients. *J Hepatol* 2002;**37**:837-842.
27. Adinolfi LE, Gambardella M, Andreana A, Tripodi MF, Utili R, Ruggiero G. Steatosis accelerates the progression of liver damage of chronic hepatitis C patients and correlates with specific HCV genotype and visceral obesity. *Hepatology* 2001;**33**:1358-1364.
28. Leandro G, Mangia A, Hui J, et al. Relationship between steatosis, inflammation, and fibrosis in chronic hepatitis C: a meta-analysis of individual patient data. *Gastroenterology* 2006;**130**:1636-1642.
29. Abenavoli L, Masarone M, Peta V, et al. Insulin resistance and liver steatosis in chronic hepatitis C infection genotype 3. *World J Gastroenterol* 2014;**20**:15233-15240.
30. Butt AA, Yan P, Bonilla H, et al; ERCHIVES (Electronically Retrieved Cohort of HCV Infected Veterans) Study Team. Effect of addition of statins to antiviral therapy in hepatitis C virus-infected persons: Results from ERCHIVES. *Hepatology* 2015;**62**:365-374.
31. Platt L, Easterbrook P, Gower E, et al. Prevalence and burden of HCV coinfection in people living with HIV: a global systematic review and meta-analysis. *Lancet Infect Dis* 2016;**16**:797-808.
32. Berenguer J, Rivero A, Jarrin I, et al. Human immunodeficiency virus/hepatitis C virus coinfection in Spain: prevalence and patient characteristics. *Open Forum Infect Dis* 2016;**3**:ofw059.
33. Lo Re V 3<sup>rd</sup>, Kallan MJ, Tate JP, et al. Hepatic decompensation in antiretroviral-treated patients co-infected with HIV and hepatitis C virus compared with hepatitis C virus-monoinfected patients: a cohort study. *Ann Intern Med* 2014;**160**:369-379.
34. Abdel-Hakeem MS, Shoukry NH. Protective immunity against hepatitis C: many shades of gray. *Front Immunol* 2014;**5**:274.
35. Konerman MA, Mehta SH, Sutcliffe CG, et al. Fibrosis progression in human immunodeficiency virus/hepatitis C virus coinfecting adults: prospective analysis of 435 liver biopsy pairs. *Hepatology* 2014;**59**:767-775.
36. Chen JY, Feeney ER, Chung RT. HCV and HIV coinfection: mechanisms and management. *Nat Rev Gastroenterol Hepatol* 2014;**11**:362-371.
37. Peters L, Klein MB. Epidemiology of hepatitis C virus in HIV-infected patients. *Curr Opin HIV AIDS* 2015;**10**:297-302.
38. Pineda JA, García-García JA, Aguilar-Guisado M, et al; Grupo para el Estudio de las Hepatitis Viricas de la Sociedad Andaluza de Enfermedades Infecciosas (SAEI). Clinical progression of hepatitis C virus-related chronic liver disease in human immunodeficiency virus-infected patients undergoing highly active antiretroviral therapy. *Hepatology* 2007;**46**:622-630.
39. Bräu N, Salvatore M, Ríos-Bedoya CF, et al. Slower fibrosis progression in HIV/HCV-coinfecting patients with successful HIV suppression using antiretroviral therapy. *J Hepatol* 2006;**44**:47-55.
40. Qurishi N, Kreuzberg C, Lüchters G, et al. Effect of antiretroviral therapy on liver-related mortality in patients with HIV and hepatitis C virus coinfection. *Lancet* 2003;**362**:1708-1713.
41. Lin W, Weinberg EM, Chung RT. Pathogenesis of accelerated fibrosis in HIV/HCV coinfection. *J Infect Dis* 2013;**207**(Suppl 1):S13-S18.
42. Berenguer J, Rodríguez-Castellano E, Carrero A, et al; GESIDA HIV/HCV Cohort Study Group. Eradication of hepatitis C virus and non-liver-related non-acquired immune deficiency syndrome-related events in human immunodeficiency virus/hepatitis C virus coinfection. *Hepatology* 2017;**66**:344-356.
43. Limketkai BN, Mehta SH, Sutcliffe CG, et al. Relationship of liver disease stage and antiviral therapy with liver-related events and death in adults coinfecting with HIV/HCV. *JAMA* 2012;**308**:370-378.
44. Kumar RN, Balba GP. Managing the HIV/HCV-co-infected patient in the direct-acting antiviral era: a review of pertinent drug interactions. *Curr Treat Options Infect Dis* 2017;**9**:411-424.
45. Douam F, Lavillette D, Cosset FL. The mechanism of HCV entry into host cells. *Prog Mol Biol Transl Sci* 2015;**129**:63-107.
46. Paul D, Madan V, Bartenschlager R. Hepatitis C virus RNA replication and assembly: living on the fat of the land. *Cell Host Microbe* 2014;**16**:569-579.
47. Aizawa Y, Seki N, Nagano T, Abe H. Chronic hepatitis C virus infection and lipoprotein metabolism. *World J Gastroenterol* 2015;**21**:10299-10313.
48. Felmlee DJ, Hafirassou ML, Lefevre M, Baumert TF, Schuster C. Hepatitis C virus, cholesterol and lipoproteins—impact for the viral life cycle and pathogenesis of liver disease. *Viruses* 2013;**5**:1292-1324.
49. Saigal S, Choudhary NS, Saraf N, et al. Genotype 3 and higher low-density lipoprotein levels are predictors of good response to treatment of recurrent hepatitis C following living donor liver transplantation. *Indian J Gastroenterol* 2015;**34**:305-309.
50. Paziienza V, Clément S, Pugnale P, et al. The hepatitis C virus core protein of genotypes 3a and 1b downregulates insulin receptor substrate 1 through genotype-specific mechanisms. *Hepatology* 2007;**45**:1164-1171.
51. Poustchi H, Negro F, Hui J, et al. Insulin resistance and response to therapy in patients infected with chronic hepatitis C virus genotypes 2 and 3. *J Hepatol* 2008;**48**:28-34.
52. D'Souza R, Sabin CA, Foster GR. Insulin resistance plays a significant role in liver fibrosis in chronic hepatitis C and in the response to antiviral therapy. *Am J Gastroenterol* 2005;**100**:1509-1515.
53. Dai CY, Huang JF, Hsieh MY, et al. Insulin resistance predicts

- response to peginterferon-alpha/ribavirin combination therapy in chronic hepatitis C patients. *J Hepatol* 2009;**50**:712-718.
54. Knobler H, Stephen M. Hepatitis C and insulin action: An intimate relationship. *World J Hepatol* 2016;**8**:131-138.
  55. Milner KL, Jenkins AB, Trenell M, et al. Eradicating hepatitis C virus ameliorates insulin resistance without change in adipose depots. *J Viral Hepat* 2014;**21**:325-332.
  56. Eslam M, Aparcero R, Kawaguchi T, et al. Meta-analysis: insulin resistance and sustained virological response in hepatitis C. *Aliment Pharmacol Ther* 2011;**34**:297-305.
  57. Fattovich G, Covolo L, Bibert S, et al; ITAHEC Study Group. IL28B polymorphisms, IP-10 and viral load predict virological response to therapy in chronic hepatitis C. *Aliment Pharmacol Ther* 2011;**33**:1162-1172.
  58. Patel S, Jinjuvadia R, Patel R, Liangpunsakul S. Insulin resistance is associated with significant liver fibrosis in chronic hepatitis C patients: A systemic review and meta-analysis. *J Clin Gastroenterol* 2016;**50**:80-84.
  59. White DL, Thrift AP, Kanwal F, Davila J, El-Serag HB. Incidence of hepatocellular carcinoma in all 50 United States, from 2000 through 2012. *Gastroenterol* 2017;**152**:812-820.
  60. Hallager S, Ladelund S, Kjaer M, et al. Hepatocellular carcinoma in patients with chronic hepatitis C and cirrhosis in Denmark: a nationwide cohort study. *J Viral Hepat* 2018;**25**:47-55.
  61. Kattakuzhy S, Levy R, Rosenthal E, Tang L, Wilson E, Kottlil S. Hepatitis C genotype 3 disease. *Hepatol Intern* 2016;**10**:861-870.
  62. McMahon BJ, Bruden D, Townshend-Bulson L, et al. Infection with hepatitis C virus genotype 3 is an independent risk factor for end-stage liver disease, hepatocellular carcinoma, and liver-related death. *Clin Gastroenterol Hepatol* 2017;**15**:431-437.
  63. Kanwal F, Kramer JR, Ilyas J, Duan Z, El-Serag HB. HCV genotype 3 is associated with an increased risk of cirrhosis and hepatocellular cancer in a national sample of U.S. Veterans with HCV. *Hepatology* 2014;**60**:98-105.
  64. Ioannou GN, Green PK, Berry K. HCV eradication induced by direct-acting antiviral agents reduces the risk of hepatocellular carcinoma. *J Hepatol* 2018;**68**:25-32.
  65. Timm J, Roggendorf M. Sequence diversity of hepatitis C virus: implications for immune control and therapy. *World J Gastroenterol* 2007;**13**:4808-4817.
  66. Fried MW. Side effects of therapy of hepatitis C and their management. *Hepatology* 2002;**36**:S237-S244.
  67. Scheel TK, Rice CM. Understanding the hepatitis C virus life cycle paves the way for highly effective therapies. *Nat Med* 2013;**19**:837-849.
  68. Pearlman BL, Traub N. Sustained virologic response to antiviral therapy for chronic hepatitis C virus infection: a cure and so much more. *Clin Infect Dis* 2011;**52**:889-900.
  69. Aghemo A, Colombo M. No time for bullies in hepatitis C virus therapy: genotype 3 conquered? *Gastroenterology* 2016;**150**:1241-1243.
  70. Pawlowsky JM. New hepatitis C therapies: the toolbox, strategies, and challenges. *Gastroenterol* 2014;**146**:1176-1192.
  71. Aghemo A, De Francesco R. New horizons in hepatitis C antiviral therapy with direct-acting antivirals. *Hepatology* 2013;**58**:428-438.
  72. Ashfaq UA, Javed T, Rehman S, Nawaz Z, Riazuddin S. An overview of HCV molecular biology, replication and immune responses. *Viral J* 2011;**8**:161.
  73. Johnson SW, Thompson DK, Raccor B. Hepatitis C virus-genotype 3: update on current and emergent therapeutic interventions. *Curr Infect Dis Rep* 2017;**19**:22.
  74. Food and Drug Administration. FDA approves Mavyret for Hepatitis C. <https://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm570038.htm> [Accessed May 1, 2018]
  75. Foster GR, Afdhal N, Roberts SK, et al; ASTRAL-3 Investigators. Sofosbuvir and velpatasvir for HCV genotype 2 and 3 infection. *N Engl J Med* 2015;**373**:2608-2617.
  76. Asselah T, Kowdley KV, Zadeikis N, et al. Efficacy of hepatitis glecaprevir/pibrentasvir for 8 or 12 weeks in patients with Hepatitis C virus genotype 2, 4, 5, or 6 infection without cirrhosis. *Clin Gastroenterol Hepatol* 2018;**16**:417-426.
  77. Wyles D, Poordad F, Wang S, et al. Glecaprevir/pibrentasvir for HCV genotype 3 patients with cirrhosis and/or prior treatment experience: a partially randomized phase III clinical trial. *Hepatology* 2017;**2**:514-523.
  78. Foster GR, Pianko S, Brown A, et al. Efficacy of sofosbuvir plus ribavirin with or without peginterferon-alfa in patients with hepatitis C virus genotype 3 infection and treatment-experienced patients with cirrhosis and hepatitis C virus genotype 2 infection. *Gastroenterology* 2015;**149**:1462-1470.
  79. AASLD/IDSA recommendations for testing, managing, and treating Hepatitis C. <https://www.hcvguidelines.org/> [Accessed May 1, 2018]
  80. Lawitz E, Lalezari JP, Hassanein T, et al. Sofosbuvir in combination with peginterferon alfa-2a and ribavirin for non-cirrhotic, treatment-naive patients with genotypes 1, 2, and 3 hepatitis C infection: a randomised, double-blind, phase 2 trial. *Lancet Infect Dis* 2013;**13**:401-408.
  81. Lawitz E, Mangia A, Wyles D, et al. Sofosbuvir for previously untreated chronic hepatitis C infection. *N Engl J Med* 2013;**368**:1878-1887.
  82. Jacobson IM, Gordon SC, Kowdley KV, et al; FUSION Study. Sofosbuvir for hepatitis C genotype 2 or 3 in patients without treatment options. *N Engl J Med* 2013;**368**:1867-1877.
  83. Younossi ZM, Stepanova M, Zeuzem S, et al. Patient-reported outcomes assessment in chronic hepatitis C treated with sofosbuvir and ribavirin: the VALENCE study. *J Hepatol* 2014;**61**:228-234.
  84. Nelson DR, Cooper JN, Lalezari JP, et al; ALLY-3 Study Team. All-oral 12-week treatment with daclatasvir plus sofosbuvir in patients with hepatitis C virus genotype 3 infection: ALLY-3 phase III study. *Hepatology* 2015;**61**:1127-1135.
  85. Leroy V, Angus P, Bronowicki JP, et al. Daclatasvir, sofosbuvir, and ribavirin for hepatitis C virus genotype 3 and advanced liver disease: A randomized phase III study (ALLY-3+). *Hepatology* 2016;**63**:1430-1441.
  86. Lawitz E, Sullivan G, Rodriguez-Torres M, et al. Exploratory trial of ombitasvir and ABT-450/r with or without ribavirin for HCV genotype 1, 2, and 3 infection. *J Infect* 2015;**70**:197-205.
  87. Gane EJ, Hyland RH, An D, et al. Sofosbuvir/ledipasvir fixed dose combination is safe and effective in difficult-to-treat populations including genotype-3 patients, decompensated genotype-1 patients, and genotype-1 patients with prior sofosbuvir treatment experience. *J Hepatol* 2014;**60**:S3-S4.
  88. Berden FA, Aaldering BR, Groenewoud H, Int'Hout J, Kievit W, Drenth JP. Identification of the best direct-acting antiviral regimen for patients with hepatitis C virus genotype 3 infection: a systematic review and network meta-analysis. *Clin Gastroenterol Hepatol* 2017;**15**:349-359.
  89. Wyles D, Bräu N, Kottlil S, et al; ASTRAL-5 Investigators. Sofosbuvir and velpatasvir for the treatment of hepatitis C virus type 1: an open-label, phase 3 study. *Clin Infect Dis* 2017;**65**:6-12.
  90. European Association for the Study of the Liver. Electronic address: [easloffice@easloffice.eu](mailto:easloffice@easloffice.eu). EASL Recommendations on Treatment of Hepatitis C 2016. *J Hepatol* 2017;**66**:153-194.
  91. Chhatwal J, Kanwal F, Roberts MS, Dunn MA. Cost-effectiveness and budget impact of hepatitis C virus treatment with sofosbuvir and ledipasvir in the United States. *Ann Intern Med* 2015;**162**:397-406.
  92. Najafzadeh M, Andersson K, Shrank WH, et al. Cost-effectiveness

- of novel regimens for the treatment of hepatitis C virus. *Ann Intern Med* 2015;**162**:407-419.
93. Ruiz I, Chevaliez S, Pawlotsky JM. Next-generation direct-acting antiviral drug-based regimens for hepatitis C. *Curr Hepatology Rep* 2017;**16**:184-191.
  94. Lawitz E, Poordad F, Gutierrez JA, et al. Short-duration treatment with elbasvir/grazoprevir and sofosbuvir for hepatitis C: a randomized trial. *Hepatology* 2017;**65**:439-450.
  95. Foster GR, Agarwal K, Cramp ME, et al. Elbasvir/Grazoprevir and Sofosbuvir for HCV genotype 3 infection with compensated cirrhosis: a randomized trial. *Hepatology* 2018;**67**:2113-2126.
  96. Kowdley KV, Lawitz E, Poordad F, et al. Phase 2b trial of interferon-free therapy for hepatitis C virus genotype 1. *N Engl J Med* 2014;**370**:222-232.
  97. Carrion AF, Gutierrez J, Martin P. New antiviral agents for the treatment of hepatitis C: ABT-450. *Expert Opin Pharmacother* 2014;**15**:711-716.
  98. Shafran SD, Shaw D, Charafeddine M, et al. Efficacy and safety results of patients with HCV genotype 2 or 3 infection treated with ombitasvir/paritaprevir/ritonavir and sofosbuvir with or without ribavirin (QUARTZ II-III). *J Viral Hepat* 2018;**25**:118-125.
  99. Gane E, Poordad F, Wang S, et al. High efficacy of ABT-493 and ABT-530 treatment in patients with HCV genotype 1 or 3 infection and compensated cirrhosis. *Gastroenterology* 2016;**151**:651-659.e1.
  100. Lawitz E, Gane EJ, Feld JJ, et al. C-BREEZE 2: efficacy and safety of a two-drug direct-acting antiviral agent (DAA) regimen ruzasvir 180 mg and uprifosbuvir 450 mg for 12 weeks in adults with chronic hepatitis C virus (HCV) genotype (GT) 1, 2, 3, 4, or 6. *Hepatology* 2017;**66**:A34-A35.
  101. Lawitz E, Buti M, Vierling JM, et al. Safety and efficacy of a fixed-dose combination regimen of grazoprevir, ruzasvir, and uprifosbuvir with or without ribavirin in participants with and without cirrhosis with chronic hepatitis C virus genotype 1, 2, or 3 infection (C-CREST-1 and C-CREST-2, part B): two randomised, phase 2, open-label trials. *Lancet Gastroenterol Hepatol* 2017;**2**:814-823.
  102. Borgia G, Maraolo AE, Nappa S, Gentile I, Buonomo AR. NS5B polymerase inhibitors in phase II clinical trials for HCV infection. *Expert Opin Investig Drugs* 2018;**27**:243-250.
  103. Foster G, Peter JR, Sergio B, et al. A randomized, phase 3 trial of sofosbuvir/velpatasvir/voxilaprevir for 8 weeks and sofosbuvir/velpatasvir for 12 weeks for patients with genotype 3 HCV infection and cirrhosis: the POLARIS-3 study. American Association for the Study of Liver Diseases Annual Meeting; 11-15 Nov 2016; Boston.
  104. Lindqvist A, Kylefjord H, Wähling H, et al. Preclinical characterization of MIV-802, a novel uridine nucleotide HCV NS5B polymerase inhibitor, for treatment of hepatitis C virus infection. *J Hepatol* 2015;**62**:S58.
  105. Lana E, Brenndorfer E, Kylefjord H, et al. MIV-802, a uridine nucleotide prodrug, is a more potent inhibitor of hepatitis C virus genotype 3 replication than sofosbuvir. *J Hepatol* 2017;**66**:S322-S323.
  106. Morisco F, Granata R, Camera S, et al; ITAL-C consortium. Optimization of direct anti-viral agent treatment schedule: Focus on HCV genotype 3. *United European Gastroenterol J* 2018;**6**:225-237.
  107. Naggie S, Muir AJ. Oral combination therapies for hepatitis C virus infection: successes, challenges, and unmet needs. *Annu Rev Med* 2017;**68**:345-358.
  108. Falade-Nwulia O, Suarez-Cuervo C, Nelson DR, Fried MW, Segal JB, Sulkowski MS. Oral direct-acting agent therapy for hepatitis C virus infection: a systematic review. *Ann Intern Med* 2017;**166**:637-648.
  109. Dore GJ, Feld JJ. Hepatitis C virus therapeutic development: in pursuit of "perfectovir". *Clin Infect Dis* 2015;**60**:1829-1836.
  110. Grebely J, Dore GJ, Morin S, Rockstroh JK, Klein MB. Elimination of HCV as a public health concern among people who inject drugs by 2030 - what will it take to get there? *J Int AIDS Soc* 2017;**20**:22146.
  111. Falade-Nwulia O, Sulkowski MS, Merkow A, Latkin C, Mehta SH. Understanding and addressing hepatitis C reinfection in the oral direct-acting antiviral era. *J Viral Hepat* 2018;**25**:220-227.
  112. Stone J, Martin NK, Hickman M, et al. The potential impact of a hepatitis c vaccine for people who inject drugs: is a vaccine needed in the age of direct-acting antivirals? *PloS One* 2016;**11**:e0156213.
  113. Liang TJ. Current progress in development of hepatitis C virus vaccines. *Nat Med* 2013;**19**:869-878.
  114. Walker CM. Designing an HCV vaccine: a unique convergence of prevention and therapy? *Curr Opin Virol* 2017;**23**:113-119.